


The Version 2 VIIRS+CrIS Fusion Radiance products

E. Eva Borbas, Elisabeth Weisz, W. Paul Menzel, Chris Moeller,
Geoff Cureton*, Greg Quinn*, and Bryan Baum**

Space Science and Engineering Center, University of Wisconsin-Madison

*NASA Atmosphere-SIPS

**Retired




MODIS/VIIRS STM, May 2-4, 2023

Goal

- The VIIRS+CrIS Fusion Radiance (FSNRAD) products have been created to provide a path for continuity of products based on the Terra, Aqua, SNPP, and NOAA-20 platforms.
- *Why is this work important?* MODIS has three channels sensitive to CO₂ in the 4.5 μm CO₂ band, four channels in the broad 15 μm CO₂ band, 2 channels sensitive to H₂O near 6.7 μm, and an ozone channel near 9 μm. VIIRS has none of these IR absorption bands. The lack of the CO₂ and H₂O channels results in a degradation of the accuracy of the cloud mask especially at night in high latitudes, other cloud products (cloud top pressure/height and thermodynamic phase) and the moisture products (total precipitable water vapor, upper tropospheric humidity).
- We addressed this restriction by constructing similar Aqua MODIS IR band radiances for VIIRS based on a fusion method that uses collocated VIIRS and CrIS data.



Imager+Sounder Spatial Fusion Schematics

Step 1



Step 2


Radiance Fusion



LORES/HIRES ... low/high spatial resolution

Imager+Sounder Radiance Fusion Example

- Imager+sounder radiance fusion applied to VIIRS+CrIS to construct missing VIIRS CO₂ and H₂O absorption bands (i.e., MODIS-like bands).
- Can be applied to various instrument pairs (e.g., AVHRR+IASI, AVHRR+HIRS, VIIRS+TROPOMI, ABI+CrIS)



VIIRS+CrIS FSNRAD Product (on full VIIRS spatial resolution)

MODIS Infrared bands		VIIRS+CrIS Fusion Infrared bands		Primary Use
band	Central Wavelength [μm]	band	Central Wavelength [μm]	
23	4.05	M13	4.05	Atmospheric temperature
24	4.47	M24 Fusion	4.47	Atmospheric temperature
25	4.52	M25 Fusion	4.52	Atmospheric temperature
27	6.72	M27 Fusion*	6.72	Water vapor
28	7.33	M28 Fusion	7.33	Water vapor
29	8.55	M14	8.55	Surface and cloud properties
30	9.73	M30 Fusion	9.73	Ozone
31	11.03	M15 M15 Fusion**	10.76	Surface and cloud properties
32	12.02	M16 M16 Fusion**	12.01	Surface and cloud properties
33	13.34	M33 Fusion*	13.34	Cloud properties
34	13.64	M34 Fusion*	13.64	Cloud properties
35	13.94	M35 Fusion*	13.94	Cloud properties
36	14.23	M36 Fusion*	14.23	Cloud properties

*FSNRAD_SS subset for the CERES team

**BT diff for M15 and M16 are also provided for uncertainty estimate

Status of the VIIRS+CrIS FSNRAD products


- **V1 released at NASA LAADS DAAC:** Fall 2019
- **V2 released at NASA LAADS DAAC:** March 8, 2022
 - DOI: 10.5067/VIIRS/FSNRAD_L2_VIIRS_CRIS_SNPP.002
- **Subsetter products are available at A-SIPS:**
 - <https://sips.ssec.wisc.edu/#/products/availability?id=14372>



Product Name	Description	Available at
FSNRAD_L2_VIIRS_CRIS_SNPP	S-NPP/VIIRS Fusion Radiances	LAADS DAAC
FSNRAD_L2_VIIRS_CRIS_NOAA20	NOAA20/VIIRS Fusion Radiances	LAADS DAAC
FSNRAD_L2_VIIRS_CRIS_SS_SNPP	S-NPP/VIIRS Subsetted Fusion Radiances	Atmosphere-SIPS
FSNRAD_L2_VIIRS_CRIS_SS_NOAA20	NOAA20/VIIRS Subsetted Fusion Radiances	Atmosphere-SIPS

- **Note for SNPP:** CrIS anomaly in LW data
 - May 21 –July 12, 2021: fill value for Band 30-36 (anomaly of CrIS LW channels)
 - July 14, 2021 - fill value for Band 27, 28, B30-36 restored (Side 1 -> Side 2)


FUSRAD BT27 [K] NOAA20 on 2020020 at 1842 V1.0.1



V1


V2 update:
Decreasing
surface effect
in WV bands

FUSRAD BT27 [K] NOAA20 on 2020020 at 1842 V2.0




V2

FUSRAD BT27 [K] NOAA20 on 2020020 at 1842 V1.0.1




FUSRAD BT27 [K] NOAA20 on 2020020 at 1842 V2.0




Aqua/MODIS B27

April 26, 2021




N20/FUS V1 B27



Aqua/MODIS B27


April 26, 2021



210

>=255



N20/FUS V2 B27



Assessment of Product Quality

SNPP and NOAA20 fusion radiance products are compared directly with Aqua/MODIS measured radiances operationally.




2022-Jan MODIS-NOAA20/VIIRS Matchup Counts




Assessment of Product Quality

SNPP and NOAA20 fusion radiance products are compared directly with Aqua/MODIS measured radiances operationally.


2022-Jan MODIS-NOAA20/VIIRS Matchup Counts



Application for the CERES team



- Operational use in CERES-NOAA20 Ed1B Production
- helps to pick up clouds over snow-ice covered surfaces at nighttime
- consistency between CERES-Aqua Ed4 and CERES-NOAA20 Ed1B is improved
- FSNRAD B27 & B33 are used in narrow-band to broad-band calculation for CERES-N20 production processing




Cloud Application

(Li et al. 2020) used the 6.7 and 13.3 μm CrIS+VIIRS fusion bands in CLAVR-X, the NOAA operational cloud processing package. They demonstrated that the fusion radiances improved cloud parameters, *like cloud mask (polar regions), type/phase, and cloud height for all latitudes.*








Bias distribution of cloud top height of ice phase clouds between S-NPP VIIRS and CALIPSO/CALIOP for emissivity range a) 0 to 0.4; b) 0.4 to 0.8; c) 0.8 to 1.0; and d) 0 to 1.0. Solid and dashed lines indicate data with/without fusion channels.

Significant improvement is found for all ice cloud emissivities but especially for semi-transparent ice clouds, when the spectral information is used what the FUSION products provide.











MOD07 TPW Application

TPW results and differences with MODIS for April 9, 2018.



MOD07 TPW Application


TPW results and differences with MODIS for April 9, 2018.



VIIRS+CrIS fusion TPW and MODIS TPW remain within 1mm for all three latitude bands (mid-latitudes north, tropics and mid-latitudes south).

(Borbas et al., 2021)


MOD07 TPW Application (cont.)



Borbias et al. (2021) show the advantage of using the VIIRS+CrIS fusion radiances for IR absorption bands at 4.5, 6.7, 7.3, 13.3, 13.6, 13.9, and 14.2 μm to determine TPW and demonstrate the potential for continuity of the Terra/Aqua MODIS infrared water vapor products. This study established the feasibility of extending the MODIS IR TPW and UTH into the future. The VIIRS+CrIS fusion TPW product, supplemented with the missing IR bands, was implemented using the same approach as the MODIS TPW product. Note that the fusion-based TPW product is in excellent agreement with MODIS.

PATMOS-X Application


PATMOS-X V6 now includes the AVHRR+HIRS radiance fusion for the multidecadal NOAA POES and EUMETSAT METOP satellite record.



Other Applications

- **GEO/LEO Application:** *spatial and temporal fusion* (between imager radiances from subsequent time steps) - (*Weisz et al., 2020*) (*Anheuser et al., 2020*)
 - ABI/CrIS fusion captures the rapidly evolving atmospheric changes during thunderstorm & tornado development (when CrIS overpass is timely) (*Smith et al, 2020*)
 - Volcanic SO₂ detection at nighttime is demonstrated with ABI/CrIS fusion (*Weisz and Menzel, 2022*)

Product Fusion Example



- “**Product fusion**” is used to transfer retrieval products at low spatial resolution (LORES) to high spatial resolution (HIRES)
 - Imager / sounder product fusion has been demonstrated for VIIRS, ABI, AHI / CrIS, TROPOMI


References:

- Foster, M. J., C. Phillips, A. K. Heidinger, E. E. Borbas, Y. Li, W. P. Menzel, A. Walther, and E. Weisz, 2023: PATMOS-x Version 6.0: 40 Years of Merged AVHRR and HIRS Global Cloud Data. *J. Climate*, **36**, 1143–1160, <https://doi.org/10.1175/JCLI-D-22-0147.1>
- Weisz, E. and W. P. Menzel, 2022: Tracking atmospheric moisture changes in convective storm environments using GEO ABI and LEO CrIS data fusion. (*Rem. Sens. Env.* 14(21), 5327; <https://doi.org/10.3390/rs14215327>)
- Weisz, E. and W. P. Menzel, 2022: Monitoring the 2021 Cumbre Vieja Volcanic Eruption Using Satellite Multi-Sensor Data Fusion. (*Jour. of Geophys. Res. Atmos.*, 128, e2022JD037926. <https://doi.org/10.1029/2022JD037926>)
- Borbas, E. E., E. Weisz, C. Moeller, W. P. Menzel, and B. A. Baum, 2021: Improvement in tropospheric moisture retrievals from VIIRS through the use of infrared absorption bands constructed from VIIRS and CrIS data fusion, *Atmospheric Measurement Techniques*, 14, 1191–1203, <https://doi.org/10.5194/amt-14-1191-2021>
- Smith Sr, William & Zhang, Qi & Shao, M. & Weisz, E.. (2020). Improved Severe Weather Forecasts Using LEO and GEO Satellite Soundings. *Journal of Atmospheric and Oceanic Technology*. 37. <http://doi.org/10.1175/JTECH-D-19-0158.1>
- Weisz, E., and W. P. Menzel, 2020: An Approach to Enhance Trace Gas Determinations through Multi-Satellite Data Fusion, *J. Appl. Remote Sens.*, 14(4), 044519, <http://doi.org/10.1117/1.JRS.14.044519>
- Anheuser, J., E. Weisz, and W. P. Menzel, 2020: Low earth orbit sounder retrieval products at geostationary earth orbit spatial and temporal scales, *J. Appl. Remote Sens.* 14(4), 048502, <http://doi.org/10.1117/1.JRS.14.048502>
- Li, Y., B. A. Baum, A. K. Heidinger, W. P. Menzel, and E. Weisz, 2020: Improvement in cloud retrievals from VIIRS through the use of infrared absorption channels constructed from VIIRS-CrIS data fusion, *Atmospheric Measurement Techniques*, 13, 4035–4049, <https://doi.org/10.5194/amt-13-4035-2020>
- Weisz, E., and W. P. Menzel, 2020: An Approach to Enhance Trace Gas Determinations through Multi-Satellite Data Fusion, *J. Appl. Remote Sens.*, 14(4), 044519 (2020), <http://doi.org/10.1117/1.JRS.14.044519>
- Weisz, E., and W. P. Menzel, 2019: Imager and sounder data fusion to generate sounder retrieval products at an improved spatial and temporal resolution, *J. Appl. Remote Sens.* 13(3), 034506, <http://doi.org/10.1117/1.JRS.13.034506>
- Weisz, E., B. A. Baum, and W. P. Menzel, 2017: Fusion of satellite-based imager and sounder data to construct supplementary high spatial resolution narrowband IR radiances. *J. Appl. Remote Sens.* 11(3), 036022, <http://doi.org/10.1117/1.JRS.11.036022>

Aqua/MODIS B33

April 26, 2021

Thank you!



NOAA-20/FSNRAD V2 B33